




Physics with Cold Molecules Using Buffer Gas Cooling

Precision Measurement, Collisions, and Laser Cooling

Nick Hutzler, Doyle Group, Harvard University


Buffer gas cooling

- › Use inert gas in cryogenic environment to cool via collisions
- › Can be used to make cold atoms or molecules in a cell or beam
- › Highly versatile – works for radicals, refractory materials, reactive species, polyatomics,...
- › Also called collisional cooling [Messer & De Lucia, 1984]
- › Review: N. Hutzler, H. Lu, & J. Doyle, Chem. Rev. 112, 4803 (2012)



Hydrodynamic Buffer Gas Beams

- › Entraining cold species in buffer flow results in high extraction (10-40%)^[1,2,3,4]
- › Translationally, rotationally cold
- › Velocity 100–200 m/s
- › >100x brighter than comparable supersonic beams^[5]




[1] S. Maxwell et al., PRL 95, 173201 (2005); [2] D. Patterson & J. Doyle, JCP 126, 154307 (2007)

[3] N. Hutzler et al., PCCP 13, 18976 (2011); [4] J. Barry et al., PCCP 13, 18936 (2011)

[5] M. Tarbutt et al., J. Phys B. 35, 5013 (2002)



ACME Electron EDM Search

- › Largest accessible electric fields are inside polar molecules
 - 10-100 GV/cm
- › Look for $d_e \cdot \mathcal{E}_{\text{eff}}$ shifts in levels
- › Molecules are typically heavy, reactive, radicals
 - Our molecule: ThO ($H^3\Delta_1$)
 - YbF, ThF⁺, WC, PbF, PbO,...
- › Slow, bright, cold beam source is desirable



ACME Results

- › $|d_e| < 8.7 \times 10^{-29} \text{ e cm}^{[1]}$
 - Ten-fold improvement^[2]
 - Shot noise limited
 - Currently working on another factor of 10
- › Theoretical and experimental study of Zeeman interaction
 - Effective g-factors depend on E-field, molecule orientation^[3]
 $\text{Energy} \propto M\Omega|E|B$
 - $\sim 10^{-5} \mu_B$ at 100 V/cm
 - Measured effect at $\sim 10^{-7} \mu_B$ level, agrees with calculations^[4]





[1] ACME Collaboration, Science 343, 269 (2014) [2] J. Hudson et al., Nature 473, 493 (2011)

[3] S. Bickman et al., PRA 80, 023418 (2009) [4] A. Petrov et al., PRA 89, 062505 (2014)

Slow Beams

- › Place second cell in front of beam aperture
 - Add gaps, vents to create low steady-state pressure
 - Cool to 1 K
- › Resulting beam is nearly-effusive, with molecules as slow as 20 m/s^[1,2]
- › ~10x less intense compared to hydrodynamic beams
- › Useful for trap loading
 - MOTs of Yb, Tm, Er, Ho without slowing^[3]



[1] D. Patterson & J. Doyle, JCP 126, 154307 (2007); [2] H. Lu et al., PCCP 42, 18986 (2011)

[3] B. Hemmerling et al., arXiv:1310.3239, to appear in New J. Phys. (2014)


Laser Cooling Polar Molecules

› Molecules lack cycling transitions

› Possible to find quasi-cycling transitions

- Choose molecules with $J=3$ diagonal Franck-Condon factors^[1]
- Use $J \rightarrow J-1$ transition^[2]

› First realized in SrF, DeMille Group @ Yale^[3] using buffer gas beam




[1] M. Di Rosa, EJP D 31, 395 (2004); [2] B. Stuhl et al., PRL 101, 243002 (2008)

[3] E. Shuman et al., Nature 467, 820 (2010)



Magneto-Optical Trapping

- › Difficulty: $J \rightarrow J-1$ transitions have dark Zeeman states
 - MOTs rely on scattering both circular polarizations
- › Our approach: change sign of B, polarization at $\sim \gamma_{\text{rad}}$
 - Molecules cannot accumulate in dark states
 - Collaboration with Ye Group @ JILA, who first demonstrated in 2D MOT with YO^[2]
- › DC MOT of SrF demonstrated recently^[1] by DeMille Group @Yale






Testbed: AC MOT on ^6Li D1 Line

- › Li D1 line has similar dark state problem ($F \geq F'$)
- › AC MOT at 6 MHz yields significant increase in number, lifetime over DC MOT
 - Will this give us an improved MOT compared to SrF?



Progress Toward CaF MOT




- › Goal: dense, long-lived MOT of CaF
- › White-light slowing
 - Recently: ~50 m/s slowing with hydrodynamic beam
 - Also demonstrated at ICL with supersonic^[1]
- › Next steps
 - Optimize slowing
 - Load MOT



[1] V. Zhelyazkova et al., PRA 89, 053416 (2014)



Magnetic Trapping of CaF and K

- › Optically pump into trapped state
 - Sufficient energy removed by climb into trap
 - General: magnetic moment, 2 photon cycles
- › Recently demonstrated with CaF^[1] and K




Results

- › Demonstrated trap lifetime of 500 ms for CaF, 9 s for K
- › Next steps
 - Co-load atom, molecule
 - Observe collisions
 - Sympathetically cool?
- › Extend to triatomics
 - Path to ultracold polyatomics?
 - Observed cold SrOH in buffer cell [Tuesday]
 - Spectroscopy from T. Steimle was a great help!



Applications with Large Molecules

- › Broadband mixture analysis
 - Cold, slow-moving, dense samples
 - Combine with optical, FTMW, double-resonance, comb, etc.
- › Enantiomer-specific detection^[1]
 - Fast, sensitive, widely applicable



[1] D. Patterson, M. Schnell, & J. M. Doyle, Nature **497**, 475 (2013)



Large Molecules

› Recent Measurements in Benzonitrile^[1]

- Direct observation of dimer formation


› When is a molecule too large to buffer gas cool?

- Clustering
- Not known
- Recently demonstrated non-sticking of Nile Red, trans-Stilbene^[2]




Fundamental Physics with Cold Molecules

Cold Chemistry and Collisions




K.-K. Ni et al., Nature **464**, 1324 (2010)

Quantum Matter/Simulation




A. Micheli et al., Nat. Phys. **2**, 341 (2006)

Quantum Computation



D. DeMille, PRL **88**, 67901 (2002)

Precision Measurement



ACME, Science **343**, 269 (2014)

Much Chemical Reviews **112** (2012)

more! L. Carr, D. DeMille, R. Krems, & J. Ye, New J. Phys. **11**, 55049 (2009)

John Doyle's Group + Collaborators

MOT

Eunmi Chae
Garrett Drayna
Börge Hemmerling
Nick Hutzler
Aakash Ravi
Wolfgang Ketterle, MIT

JILA Collaborators

Alejandra Collopy
Matthew Hummon
Mark Yeo
Yong Xia
Jun Ye

ACME EDM

Doyle Group

Jacob Baron
Nick Hutzler
Elizabeth Petrik
Gabrielse Group
Paul Hess (→JQI)
Cris Panda
Ben Spaun (→JILA)

DeMille Group @ Yale

Brendon O'Leary
Adam West

Collaborators

Wes Campbell, UCLA
Amar Vutha, York

Theory @ PNPI

Alexander Petrov
Leonid Skripnikov
Anatoly Titov

Large Molecules

Dave Patterson
Julia Piskorski
Sandra Eibenberger, Vienna
Melanie Schnell, CFEL

Optical Loading

Louis Baum
Börge Hemmerling
Ivan Kozyryev
Hsin-I Lu (→JQI)
Julia Piskorski

Funding

NSF, DOE, AFOSR MURI

Review:

N. Hutzler, H. Lu, & J.
Doyle, Chem. Rev. 112,
4803 (2012)

Harvard/MIT Center for Ultracold Atoms (CUA)

www.doylegroup.harvard.edu

